

| Lake Characteristics | | | | | |
|--------------------------|---|--|--|--|--|
| Surface Area (ac/ha) | 659/267 | | | | |
| Max Depth (ft/m) | 24.6/7.5 | | | | |
| Mean Depth (ft/m) | 12.1/3.7 | | | | |
| Retention Time (years) | 0.69 | | | | |
| Lake Classification | А | | | | |
| Dam Classification | А | | | | |
| Watershed Characteristic | s | | | | |
| Watershed area (ac/ha) | 7562/3060 | | | | |
| Watershed/Lake Ratio | 11 | | | | |
| Lake and Wetlands % | 17.04% | | | | |
| Agricultural % | 20.72% | | | | |
| Forest, Shrub, Grasses % | 56.61% | | | | |
| Residential % | 5.62% | | | | |
| Urban % | 0.01% | | | | |
| CSLAP Participation | | | | | |
| Years in CSLAP | 1992-1996/ 1998-1999/ 2001-2010/ 2012-2021 | | | | |
| Volunteers | Lynn Clauer, Lynn Wilbur, bill Zeppetelli | | | | |

| Trophic State | HABs Susceptibility | Invasive Vulnerability | PWL Assessment |
|---------------|--|---------------------------------------|--|
| Eutrophic | No Reported Blooms, Moderate susceptibility | Invasives present, High vulnerability | Download Assessment Here (https://www.dec.ny.gov/data/W 0002.html) |

Cossayuna Lake – 2021 Sampling Season Results

Red shaded results indicate eutrophic water quality conditions. In these columns, 'No' indicates there was no significant trend, ' \uparrow ' indicates there was a significant positive trend (p<0.01), ' \downarrow ' indicates there was a significant positive trend (p<0.01), ' \downarrow ' indicates there was a significant negative trend (p<0.01), and blank indicates there was insufficient data to identify a trend. In this report, seasonal trend analyses for individual sampling years and long term trend analyses show changes in key water quality indicators over a consistent index period (mid-June thru mid-September).

| 2021 Sampling Results | | | | | | | | | | | | | | |
|--------------------------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|--------------------|------------------|-------------------|------------------|--------------------|-------------------|
| Open Water Indicators | 06- 01 | 06- 15 | 06- 29 | 07- 13 | 07- 27 | 08- 10 | 08- 23 | 09- 07 | Seasonal Change | Season Median | Decadal Median | Decadal Trend | Longterm Median | Longterm Trend |
| Clarity (m) | 5 | 4.2 | 2.8 | 2.6 | 2.9 | 2.1 | 2 | 1.5 | <u></u> | 2.7 | 2.6 | no | 1.8 | 11 |
| Deep Temp (degC) | 16 | | | 20 | | 25 | | 20 | <u></u> | 20 | 22 | no | 22 | no |
| Upper Temp (degC) | 17 | | | 24 | | 27 | | 25 | | 24.5 | 24 | no | 23 | 11 |
| Surface TP (mg/L) | 0.034 | 0.016 | 0.023 | | | 0.021 | 0.022 | 0.025 | \ | 0.022 | 0.022 | no | 0.027 | 11 |
| Surface TDP (mg/L) | 0.015 | 0.027 | 0.01 | | | 0.014 | 0.007 | | ^ | 0.014 | 0.01 | no | 0.01 | no |
| Deep TP (mg/L) | 0.024 | 0.017 | 0.032 | 0.046 | | 0.036 | 0.212 | 0.056 | | 0.036 | 0.027 | no | 0.029 | 1 |
| Deep TDP (mg/L) | 0.02 | 0.01 | 0.012 | 0.02 | | 0.021 | 0.035 | 0.042 | | 0.02 | 0.02 | no | 0.02 | no |
| TN (mg/L) | 0.248 | 0.288 | | 0.317 | 0.357 | 0.446 | 0.44 | 0.598 | | 0.357 | 0.476 | 11 | 0.473 | no |
| TDN (mg/L) | 0.277 | 0.277 | 0.312 | 0.27 | 0.267 | 0.287 | 0.335 | 0.314 | ~^^ | 0.282 | 0.339 | 1 | 0.354 | 11 |
| TN:TP | 7.2 | 18.5 | | | | 20.8 | 20.4 | 24 | | 20.4 | 21.8 | no | 20.3 | 11 |
| Surface NH3 (mg/L) | 0.045 | 0.049 | | 0 | 0.069 | 0 | 0 | 0.061 | ~\\\ | 0.045 | 0.029 | no | 0.031 | no |
| Deep NH3 (mg/L) | 0.058 | 0.051 | 0 | | 0.132 | 0.062 | | | ~_^ | 0.058 | 0.048 | 11 | 0.045 | no |
| Chl.a (ug/L) | 1 | 8.9 | 12.8 | 12.6 | 11.1 | 20.3 | 22.8 | 27 | | 12.7 | 11.2 | no | 13.5 | 11 |
| рН | 8 | 8.2 | 8.4 | 8 | 8.7 | 9 | 8.9 | 7.8 | ~ | 8.3 | 7.9 | 11 | 8.1 | 11 |
| Surface Calcium (mg/L) | 23.9 | | | | 22 | | | | | 23 | 21 | no | 23.5 | no |
| Surface Chloride (mg/L) | 17 | | 18.4 | | | | | | | 17.7 | 19 | no | 19 | no |
| True Color (ptu) | 9 | 7 | 6 | 8 | 6 | 9 | 11 | 14 | ~ / | 8.5 | 8 | 1 | 10 | no |
| Cond (uS/cm) | 268.3 | 260.9 | 263.6 | 254.3 | 261.8 | 245 | 225.3 | 230.9 | ~~ | 257.6 | 197.3 | 11 | 194 | no |
| FP BG Chl.a (ug/L) | 0.5 | 0.7 | 0.5 | 1 | 1.9 | 6.3 | 13.4 | 25.8 | | 1.4 | 2 | no | 2 | no |

Cossayuna Lake - Lake Scorecard

| Water Quality Indicators | Average Year | 2021 | | |
|--------------------------------|--------------|-------------|--|--|
| Phosphorus | Eutrophic | Eutrophic | | |
| Chlorophyll A | Eutrophic | Eutrophic | | |
| Secchi | Eutrophic | Mesotrophic | | |
| Lake Perception | Poor | Fair | | |
| Shoreline Harmful Algal Blooms | Poor | Good | | |
| Open Water Algae Levels | Poor | Poor | | |
| Aquatic Invasive Species | Present | | | |

Cossayuna Lake - 2021 Lake Summary

Q. What is the condition of the lake?

A. Cossayuna Lake continues to be eutrophic, or highly productive, based on low water clarity, high algae levels (chlorophyll a), and high nutrient (phosphorous) levels. Soluble nutrients were analyzed in 2021. Some of the phosphorus in the lake is soluble, indicating some potential for more algae growth. Most of the nitrogen in the lake is soluble. The waterbody is highly alkaline or basic, with hard water, low water color, and moderately low nitrogen levels.

Q. How did this year compare to previous years?

A. Compared to previous years, conductivity and calcium were higher in 2021. Compared to previous years, chloride was lower in 2021. Water clarity (secchi), chlorophyll a, total phosphorus, color, pH, surface water temperature, deep water temperature, water quality evaluation, aquatic plant coverage and recreational evaluation in 2021 were similar to previous years.

Q. How does this lake compare to other New York lakes?

A. Compared to other New York lakes, this lake usually has higher calcium and deep water temperature. Compared to other New York lakes, this lake usually has lower chloride and more favorable aquatic plant coverage.

Q. Are there any (statistically significant) trends?

A. Over the past 30 years, clarity, surface water temperature and TN:TP have increased significantly. Over the past 30 years, chlorophyll a, deep total phosphorus, pH, surface total phosphorus and total dissolved nitrogen have decreased significantly. Over the past ten years, conductivity, deep NH3 and pH have increased significantly. Over the past ten years, total dissolved nitrogen, total nitrogen and true color have decreased significantly.

Q. Has the lake experienced open water or shoreline harmful algal blooms (HABs)?

A. Water quality conditions generally indicate a moderate susceptibility to blooms, with frequent blooms along the shoreline or in the open water.

The open water algal community in the lake is usually comprised of high cyanobacteria levels. This community is dominated by Dolichospermum. Typically, overall open water algae levels are intermediate. Overall open water toxin levels are consistently below recreational levels of concern.

This year, overall algae levels were high, with diatoms the most common taxa in open water samples, and with low cyanobacteria levels. Open water toxin levels were at times low but detectable this year.

Shoreline blooms were not reported and/or sampled this year.

Q. Have any aquatic invasive species (AIS) been reported?

A. Invasive species have been reported in this waterbody. Aquatic invasive plant and/or animal species reported include: zebra mussels, Eurasian watermilfoil, Curly leafed pondweed, water chestnut, brittle naiad. This waterbody has high vulnerability for introduction of new invasive species due to invasive species already being present. This waterbody has moderate vulnerability for establishment of invasive bivalves based on calcium levels. For more information about invasive species in the area, or to report an invasive species observation, visit NY iMapInvasives at https://www.nyimapinvasives.org/ (https://www.nyimapinvasives.org/).


Cossayuna Lake 2021 NYHABs notifications

Were there any reported HABs this season? No.

Shoreline HAB Samples 2021

There were no shoreline HAB samples taken this season.

Open Water Algae




Shoreline Algae


There is no shoreline algae or shoreline microcystin data to display from this year.

Cossayuna Lake - In-Season Trend Analysis


In Season Water Clarity




In Season Water Temperature




Lake Perception- Clarity




Lake Perception - Plants





Lake Perception - Recreation




Cossayuna Lake Long-Term Trend Analysis









Cossayuna Lake Depth Profiles

There is no depth profile data for this site.

Water Quality Assessments

The Waterbody Inventory/Priority Waterbodies List (WI/PWL) is a statewide inventory of New York's water resources that is used to track a waters ability to support its' best use(s), identify pollutant(s) causing impairment of best use(s), and follow the status of restoration, protection and other water quality activities and efforts. Data collected through CSLAP contributes to the WI/PWL. In order to be included as an assessment unit in the WI/PWL, a lake, pond, or reservoir must be at least 6.4 acres in size.

Download Lake Assessment Here (https://www.dec.ny.gov/data/WQP/PWL/1103-0002.html)

Total Maximum Daily Load (TMDL)

A TMDL is a type of regulatory clean water plan that calculates the maximum amount of a single pollutant that a waterbody can receive and still meet water quality standards. Section 303(d) of the Clean Water Act also requires states to identify impaired waters, where designated uses are not fully supported. These waterbodies are then listed on the Clean Water Act 303(d) "impaired waters" list. Waterbodies may have been identified as impaired due to fish consumption advisories, shellfishing closures, public bathing beach closures, or sampling results (high nutrient levels, turbidity, toxic sediments). The Clean Water Act also requires states to develop TMDLs for impaired waterbodies on the 303(d) list to reduce the amount of pollutants entering impaired waterbodies to meet water quality standards. TMDL plans may also be developed to protect waterbodies from becoming impaired - for example, protecting public drinking water supplies to protect human health. DEC develops TMDLs and EPA approves them.

Download TMDL Here (https://www.dec.ny.gov/docs/water_pdf/tmdlcossayuna08.pdf)

Lake Stewardship Actions

Individual stewardship activities can help improve water quality: maintain your septic system, reduce fertilizer use, grow a buffer of native plants next to the lake shore, and reduce shoreline erosion and runoff into the lake. Visiting boats should be inspected to prevent the spread of invasive species, and continued community education about and monitoring for invasive species is recommended. Routine education about algae and harmful algal blooms (HABs) within your lake community is recommended; to learn more about HABs and see examples of HABs visit http://www.dec.ny.gov/chemical/81962.html (http://www.dec.ny.gov/chemical/81962.html). Occurrences of HABs can be reported to NYSDEC. For more information on keeping New York waters clean, visit http://www.dec.ny.gov/public/43661.html (http://www.dec.ny.gov/public/43661.html).

How to Read the Report

This guide provides a description of the CSLAP report by section and a glossary. The sampling site is indicated in the header for lakes with more than one routine sampling site.

Physical Characteristics influence lake quality:

- · Surface area is the lake's surface in acres and hectares.
- Max depth is the water depth measured at the deepest part of the lake in feet and meters.
- Mean depth is either known from lake bathymetry or is 0.46 of the maximum depth.
- Retention time is the time it takes for water to pass through a lake in years. This indicates the influence of the watershed on lake conditions.
- Lake classification describes the "best uses" for this lake. Class AA, AAspec, and A lakes may be used as sources of potable water. Class B lakes are suitable for contact recreational activities, like swimming. Class C lakes are suitable for non-contact recreational activities, including fishing, although they may still support swimming. The addition of a T or TS to any of these classes indicates the ability of a lake to support trout populations and/or trout spawning.
- Dam classification defines the hazard class of a dam. Class A, B, C, and D dams are defined as low, intermediate, high, or negligible/no hazard dams in that order. "0" indicates that no class has been assigned to a particular dam, or that no dam exists.

Watershed characteristics influence lake water quality:

- Watershed area in acres and hectares
- Land use data come from the most recent (2011) US Geological Survey National Land Use Cover dataset

CSLAP Participation lists the sampling years and the current year volunteers.

Key lake status indicators summarize lake conditions:

- Trophic state of a lake refers to its nutrient loading and productivity, measured by phosphorus, algae, and clarity. An oligotrophic lake has low nutrient and algae levels (low productivity) and high clarity while a eutrophic lake has high nutrient and algae levels (high productivity) and low clarity. Mesotrophic lakes fall in the middle.
- Harmful algal bloom susceptibility summarizes the available historical HAB data and indicates the potential for future HAB events.
- Invasive vulnerability indicates whether aquatic invasive species are found in this lake or in nearby lakes, indicating the potential for further introductions.
- Priority waterbody list (PWL) assessment is based on the assessment of use categories and summarized as fully supported, threatened, stressed, impaired, or precluded. Aesthetics and habitat are evaluated as good, fair, or poor. The cited PWL assessment reflects the "worst" assessment for the lake.

Current year sampling results shows results for each of the sampling sessions in the year are in tabular form. The seasonal change graphically shows the current year results. Red shading indicates eutrophic readings.

• If there are more than ten shoreline bloom samples collected in a year, bloom sample information is instead summarized by month (May-Oct.) as minimum, average, and maximum values for blue-green algae and microcystin.

The Lake Scorecard represents key water quality indicator results for this lake in an easy-to-read format, comparing information from the current year and historical average of the CSLAP data. Indicators include (1) trophic status of phosphorus, chlorophyll (or algae) and secchi (or clarity); (2) presence or absence of aquatic invasive plants or animals; (3) lake user perception based on perceived physical condition and recreational suitability of the lake; (4) harmful algal bloom samples or reports; and (5) algae levels in the open water of routinely sampled sites.

The Lake Summary reviews and encapsulates the data in the lake report, including comparisons to historical data from this lake, and results from nearby lakes.

Harmful Algal Blooms:

- HAB notification periods on the DEC website http://www.dec.ny.gov/chemical/83310.html (http://www.dec.ny.gov/chemical/83310.html)
- Shoreline HAB sample dates and results. Samples are collected from the area that appears to have the worst bloom. Red shading indicates
 a confirmed HAB.
- HAB sample algae analysis. Algae types typically change during the season. These charts show the amount of the different types of algae
 found in each mid-lake or shoreline sample. Samples with high levels of BGA are HABs. The second set of charts show the level of toxins
 found in open water and shoreline samples compared to NYSDOH and NYSDEC guidelines.

In-Season Trend Analysis shows water temperature and water clarity during the sampling season. These indicate seasonal changes and show the sample year results compared to the typical historical readings for those dates.

Long-Term Trend Analysis puts the current year findings in context. Summer averages (mid-June thru mid-September) for each of the CSLAP years show trends in key water quality indicators. The graphs include relevant criteria (trophic categories, water quality standards, etc.) and boundaries separating these criteria.

Glossary of Water Quality and HAB Indicators

Clarity (m): The depth to which a Secchi disk lowered into the water is visible, measured in meters. Water clarity is one of the trophic indicators for each lake.

TP (mg/L): Total phosphorus, measured in milligrams per liter at the lake surface (1.5 meters below the surface). TP includes all dissolved and particulate forms of phosphorus.

Deep TP: Total phosphorus measured in milligrams per liter at depth (1-2 meters above the lake bottom at the deepest part of the lake or a fixed depth in the hypolimnion of very deep lakes).

TN: Total nitrogen, measured in milligrams per liter at the lake surface. TN includes all forms of nitrogen, including NOx (nitrite and nitrate) and NH4 (ammonia).

N:P Ratio: The ratio of total nitrogen to total phosphorus, unitless (mass ratio). This ratio helps determine if a lake is phosphorous or nitrogen limited

Chl.a (µg/L): Chlorophyll a, measured in micrograms per liter. Indicates the amount of algae in the water column. This is an extracted chlorophyll measurement.

pH: A range from 0 to 14, with 0 being the most acidic and 14 being the most basic or alkaline. A healthy lake generally ranges between 6.5 and 8.5

Cond (µmho/cm): Specific conductance is a measure of the conductivity of water. A higher value indicates the presence of more dissolved ions. High ion concentrations (> 250) usually indicate hardwater, and low readings (< 125) usually show softwater.

Calcium (mg/L): Calcium, a component of lake buffering capacity (the ability to neutralize acid inputs), as measured in milligrams per liter at the lake surface (1.5 meters below the surface).

Chloride (mg/L): Chloride, or chloride ions, as measured in milligrams per liter at the lake surface (1.5 meters below the surface).

Upper Temp (°C): Surface temperature, measured in degrees Celsius.

Deep Temp (°C): Deep water temperature, measured in degrees Celsius.

BG Chl.a (μg/L): Chlorophyll a from blue-green algae, measured in micrograms per liter. This is an "unextracted" estimate using a fluoroprobe. This result is different from the extracted chlorophyll measurement described above.

HABs: Harmful Algal Blooms. Algal blooms that have the appearance of cyanobacteria (BGA).

BGA: Blue-green algae, also known as cyanobacteria.

Microcystin (μg/L): The most common HAB liver toxin; total microcystin above 20 micrograms per liter indicates a "high toxin" bloom. However, ALL BGA blooms pose a potential health risk and should be avoided.

Sampling Season Results Information:

Seasonal Change: Shows the current year variability

Season Median: The middle value(or average of the middle two values) of the current year's data

Decadal Median: The median of the most recent ten years of water quality data

Longterm Median: The median of all years of water quality data

Decadal & Longterm Trends: Indicate whether there was a statistically significant change in water quality data over the most recent or all years

Download Water Column Data

Download HAB Sample Data

Download Field Data